

# Concurrent Comorbidities with T2D




## Among Adults with Diabetes, 90% Experience Overweight or Obesity

Overweight (BMI 25.0-29.9 kg/m<sup>2</sup>)

Obesity (BMI 30.0-39.9 kg/m<sup>2</sup>)

Extreme obesity (BMI 40.0 kg/m<sup>2</sup> +)



## Obesity is Linked to Additional Concurrent Comorbidities

~75% with hypertension

~33-38% with PCOS

~50-75% with NAFLD

>90% with T2D



~45% with OSA

~72% with GERD


~60% with osteoarthritis

~60-70% with dyslipidemia

# Concurrent Comorbidities with T2D



## Cardiovascular Risk in T2D



Karjoo S, et al. Obesity Pillars. 2022; Barber TM, Franks S. Clin Endocrinol. 2021; Hall JE, et al. Cir Res. 2015; Bramante CT, et al. Diabetes Spectr. 2017; Feingold KR. Endotext. 2020; <https://my.clevelandclinic.org/-/scassets/files/org/bariatric/co-morbidities/gerd-and-obesity.ashx?la=en>; <https://www.obesityaction.org/wp-content/uploads/Obesity-and-Osteoarthritis-Fact-Sheet.pdf>; <https://obesitymedicine.org/obesity-and-sleep-apnea/#:~:text=Sleep%20apnea%20is%20believed%20to,45%25%20of%20individuals%20with%20obesity.>