

Shedding Light on Type 2 Diabetes: Weight Management & Future Breakthroughs



Select Bibliography and Suggested Readings

- Ahmann AJ, Capehorn M, Charpentier G, et al. Efficacy and Safety of Once-Weekly Semaglutide Versus Exenatide ER in Subjects With Type 2 Diabetes (SUSTAIN 3): A 56-Week, Open-Label, Randomized Clinical Trial. *Diabetes Care*. 2018;41(2):258-266. doi:10.2337/dc17-0417
- Ahrén B, Masmiquel L, Kumar H, et al. Efficacy and safety of once-weekly semaglutide versus once-daily sitagliptin as an add-on to metformin, thiazolidinediones, or both, in patients with type 2 diabetes (SUSTAIN 2): a 56-week, double-blind, phase 3a, randomised trial. *Lancet Diabetes Endocrinol*. 2017;5(5):341-354. doi:10.1016/S2213-8587(17)30092-X
- Aroda VR, Ahmann A, Cariou B, et al. Comparative efficacy, safety, and cardiovascular outcomes with once-weekly subcutaneous semaglutide in the treatment of type 2 diabetes: Insights from the SUSTAIN 1-7 trials. *Diabetes Metab*. 2019;45(5):409-418. doi:10.1016/j.diabet.2018.12.001.
- Aroda VR, Bain SC, Cariou B, et al. Efficacy and safety of once-weekly semaglutide versus once-daily insulin glargine as add-on to metformin (with or without sulfonylureas) in insulin-naive patients with type 2 diabetes (SUSTAIN 4): a randomised, open-label, parallel-group, multicentre, multinational, phase 3a trial. *Lancet Diabetes Endocrinol*. 2017;5(5):355-366. doi:10.1016/S2213-8587(17)30085-2
- Centers for Disease Control and Prevention. National Diabetes Prevention Program. <https://www.cdc.gov/diabetes/prevention/index.html>
- Dahl D, Onishi Y, Norwood P, et al. Effect of Subcutaneous Tirzepatide vs Placebo Added to Titrated Insulin Glargine on Glycemic Control in Patients With Type 2 Diabetes: The SURPASS-5 Randomized Clinical Trial. *JAMA*. 2022;327(6):534-545. doi:10.1001/jama.2022.0078
- Del Prato S, Kahn SE, Pavo I, et al. Tirzepatide versus insulin glargine in type 2 diabetes and increased cardiovascular risk (SURPASS-4): a randomised, open-label, parallel-group, multicentre, phase 3 trial. *Lancet*. 2021;398(10313):1811-1824. doi:10.1016/S0140-6736(21)02188-7
- Frías JP, Davies MJ, Rosenstock J, et al. Tirzepatide versus Semaglutide Once Weekly in Patients with Type 2 Diabetes. *N Engl J Med*. 2021;385(6):503-515. doi:10.1056/NEJMoa2107519
- Garvey WT, Frias JP, Jastreboff AM, et al. Tirzepatide once weekly for the treatment of obesity in people with type 2 diabetes (SURMOUNT-2): a double-blind, randomised, multicentre, placebo-controlled, phase 3 trial. *Lancet*. 2023;402(10402):613-626. doi:10.1016/S0140-6736(23)01200-X

Shedding Light on Type 2 Diabetes: Weight Management & Future Breakthroughs



- Gastaldelli A, Cusi K, Fernández Landó L, Bray R, Brouwers B, Rodríguez Á. Effect of tirzepatide versus insulin degludec on liver fat content and abdominal adipose tissue in people with type 2 diabetes (SURPASS-3 MRI): a substudy of the randomised, open-label, parallel-group, phase 3 SURPASS-3 trial. *Lancet Diabetes Endocrinol.* 2022;10(6):393-406. doi:10.1016/S2213-8587(22)00070-5
- Huang Z, Zhuang X, Huang R, et al. Physical Activity and Weight Loss Among Adults With Type 2 Diabetes and Overweight or Obesity: A Post Hoc Analysis of the Look AHEAD Trial. *JAMA Netw Open.* 2024;7(2):e240219. doi:10.1001/jamanetworkopen.2024.0219
- Lincoff AM, Brown-Frandsen K, Colhoun HM, et al. Semaglutide and Cardiovascular Outcomes in Obesity without Diabetes. *N Engl J Med.* 2023;389(24):2221-2232. doi:10.1056/NEJMoa2307563
- Ludvik B, Giorgino F, Jódar E, et al. Once-weekly tirzepatide versus once-daily insulin degludec as add-on to metformin with or without SGLT2 inhibitors in patients with type 2 diabetes (SURPASS-3): a randomised, open-label, parallel-group, phase 3 trial. *Lancet.* 2021;398(10300):583-598. doi:10.1016/S0140-6736(21)01443-4
- Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* 2016;375(4):311-322. doi:10.1056/NEJMoa1603827
- Newsome PN, Buchholtz K, Cusi K, et al. A Placebo-Controlled Trial of Subcutaneous Semaglutide in Nonalcoholic Steatohepatitis. *N Engl J Med.* 2021;384(12):1113-1124. doi:10.1056/NEJMoa2028395
- Rodbard HW, Lingvay I, Reed J, et al. Semaglutide Added to Basal Insulin in Type 2 Diabetes (SUSTAIN 5): A Randomized, Controlled Trial. *J Clin Endocrinol Metab.* 2018;103(6):2291-2301. doi:10.1210/jc.2018-00070
- Rosenstock J, Wysham C, Frías JP, et al. Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): a double-blind, randomised, phase 3 trial [published correction appears in *Lancet.* 2021 Jul 17;398(10296):212]. *Lancet.* 2021;398(10295):143-155. doi:10.1016/S0140-6736(21)01324-6
- Sattar N, McGuire DK, Pavo I, et al. Tirzepatide cardiovascular event risk assessment: a pre-specified meta-analysis. *Nat Med.* 2022;28(3):591-598. doi:10.1038/s41591-022-01707-4
- Sorli C, Harashima SI, Tsoukas GM, et al. Efficacy and safety of once-weekly semaglutide monotherapy versus placebo in patients with type 2 diabetes (SUSTAIN 1): a double-blind, randomised, placebo-controlled, parallel-group, multinational, multicentre phase 3a trial. *Lancet Diabetes Endocrinol.* 2017;5(4):251-260. doi:10.1016/S2213-8587(17)30013-X
- Wharton S, Calanna S, Davies M, et al. Gastrointestinal tolerability of once-weekly semaglutide 2.4 mg in adults with overweight or obesity, and the relationship between gastrointestinal adverse events and weight loss. *Diabetes Obes Metab.* 2022;24(1):94-105. doi:10.1111/dom.14551